Geometric Condition for Local Finiteness of a Lattice of Convex Sets

MATT INSALL

ABSTRACT. Convex sets of states and the corresponding normal functionals defined on the Hilbert space containing them play a central role in the foundations of General Quantum Mechanics (see [4, 1]). The design of quantum logics for reasoning about General Quantum Mechanics leads naturally to the theory of orthomodular lattices [4, 1, 2], for example, as lattices of closed convex sets in a (complex) Hilbert space. The quantum logics developed for such a purpose can be much easier to work in if the corresponding lattice of convex sets is finite, or at least locally finite (in the classical or intuitionistic case, this holds, because truth assignments are made in a boolean algebra a distributive lattice). We give elementary examples which establish the following facts: the lattice of convex subsets of a Hilbert space is not locally finite, it is not modular (hence not distributive), and locally finite lattices of closed convex sets in any Hilbert space have very restrictive geometric arrangements of their members.

References

- E.G. Beltrametti and G. Cassinelli, *The Logic of Quantum MEchanics*, Encyclopedia of Mathematics and its Applications, REading, Mass., Addison-Wesley, 1981.
- [2] G. Kalmbach, Orthomodular Lattices, London, Academic Press.
- [3] M.D. MacLAren, Nearly Modular Orthocomplemented Lattices, Trans. Amer. Math. Soc., 114, 401–416.
- [4] J.C.T. Poll, Orthomodular LAttices and Baer *-Semigroups in General Quantum Mechanics, Symposium on Lattice Theory and General Quantum MEchanics, Department of Mathematics, University of Missouri-Rolla, Rolla, MO, 1968.

DEPARTMENT OF MATHEMATICS AND STATISTICS UNIVERSITY OF MISSOURI-ROLLA ROLLA, MISSOURI 65401-0249 USA *E-mail address*: insall@umr.edu

¹⁹⁹¹ Mathematics Subject Classification. Primary: 06A23, 06C99; Secondary: 51M15, 52A10. Key words and phrases. Distributive lattice, modular lattice, locally finite lattice, freely generated lattice, convex sets.