

## Existence of solutions of a class of fractional partial integrodifferential equations

K. BALACHANDRAN  0000-0002-8834-7521

ABSTRACT. In this paper we study the existence of solutions for the nonlinear Sobolev type fractional partial integrodifferential equations with Neumann boundary condition. Under suitable assumptions the results are established by using the Leray-Schauder fixed point theorem.

### 1. INTRODUCTION

There has been an increasing interest in recent years shown by many researchers from various fields of science and engineering to study fractional order partial differential equations [18, 21, 24, 25, 28, 29]. These equations are found to be an effective tool to describe certain physical phenomena, such as diffusion processes [19] and viscoelasticity theories [20]. Several authors have discussed the existence of solutions for different types of fractional partial differential equations [1–3, 15, 27].

Brill [14] and Showalter [30] investigated the problem of existence of solutions of for semilinear Sobolev-type equations in Banach spaces. The Sobolev-type semilinear differential equation serves as an abstract formulation of partial differential equations which arise in various applications such as in the flow of fluid through fissured rocks [13], thermodynamics and shear in second order fluids. Lightbourne and Rankin [26] discussed the Cauchy problem for a partial functional differential equation of Sobolev-type. Balachandran et al. [12] established the existence results for Sobolev-type semilinear integrodifferential equations whereas Balachandran and Uchiyama [10] studied the same problem for nonlinear integrodifferential equation of Sobolev-type with nonlocal condition in Banach spaces. Several authors have studied the nonlocal Cauchy problem for Sobolev-type differential equations in Banach spaces [7–9, 11].

Balachandran and Kiruthika [6] investigated the existence problem for abstract fractional integrodifferential equations of Sobolev-type. Existence

---

2020 *Mathematics Subject Classification*. Primary: 35R11, 35S10; Secondary: 26A33.

*Key words and phrases*. Fractional partial integrodifferential equations, Sobolev-type, Existence of solutions.

*Full paper*. Received 21 March 2026, accepted 20 May 2026, available online 10 Jun 2026.

of solutions for abstract fractional integrodifferential equations of Sobolev-type with deviating arguments are studied in [22]. The existence results for fractional order Sobolev-type partial differential equations was discussed in [5]. In this paper, we study the existence problem for fractional partial integrodifferential equations of Sobolev-type with Neumann boundary condition by suitably adopting the technique of [27].

## 2. PRELIMINARIES

In this section, we introduce some notations and basic facts of fractional calculus. Let  $\Omega \subset \mathbb{R}$  and  $C(J, \mathbb{R})$  is the Banach space of all continuous functions from  $J = [0, T]$  into  $\mathbb{R}$ . Let  $\Gamma(\cdot)$  denote the gamma function. For any positive number  $0 < \alpha < 1$ , the Riemann Liouville derivative and Caputo derivative are defined as follows.

**Definition 1** ([23]). The Riemann-Liouville partial fractional integral operator of order  $0 < \alpha < 1$  with respect to  $t$  of a function  $z(x, t)$  is defined by

$$I^\alpha z(x, t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} z(x, s) \, ds.$$

**Definition 2** ([23]). The Riemann-Liouville partial fractional derivative of order  $0 < \alpha < 1$  of a function  $z(x, t)$  with respect to  $t$  is defined by

$$\frac{\partial^\alpha}{\partial t^\alpha} z(x, t) = \frac{1}{\Gamma(1-\alpha)} \frac{\partial}{\partial t} \int_0^t \frac{z(x, s)}{(t-s)^\alpha} \, ds.$$

**Definition 3** ([23]). The Caputo partial fractional derivative of order  $0 < \alpha < 1$  with respect to  $t$  of a function  $z(x, t)$  is defined as

$$\frac{C\partial^\alpha}{\partial t^\alpha} z(x, t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{1}{(t-s)^\alpha} \frac{\partial z(x, s)}{\partial s} \, ds.$$

A significant research development in ordinary and partial differential equations involving both Riemann-Liouville and Caputo fractional derivatives is contained in [4, 5, 27]. The Riemann-Liouville and Caputo fractional derivatives are linked by the relationship

$$\frac{C\partial^\alpha}{\partial t^\alpha} z(x, t) = \frac{\partial^\alpha}{\partial t^\alpha} z(x, t) - \frac{z(x, 0)}{\Gamma(1-\alpha)t^\alpha}.$$

Further the property

$$(1) \quad I^\alpha \frac{C\partial^\alpha z(x, t)}{\partial t^\alpha} = z(x, t) - z(x, 0)$$

is useful in proving the existence results.

**Theorem 1** (Leray-Schauder fixed point theorem, see [17]). *If  $U$  is a closed bounded convex subset of a Banach space  $X$  and  $P : U \rightarrow U$  is completely continuous, then  $P$  has at least a fixed point in  $U$ .*

**Theorem 2** (Green's Identity, see [16]). *Let  $\Omega$  be a bounded domain in  $\mathbb{R}^m$  with smooth boundary  $\partial\Omega$ . Then, for any  $u, v \in C^2(\Omega)$ ,*

$$\int_{\Omega} v \Delta u \, dx = \int_{\partial\Omega} v \frac{\partial u}{\partial n} \, ds - \int_{\Omega} \nabla u \cdot \nabla v \, dx,$$

where  $n$  is the outward unit normal to the boundary  $\partial\Omega$  and  $ds$  is the element of arc length. For the special case  $v = 1$ ,

$$(2) \quad \int_{\Omega} \Delta u \, dx = \int_{\partial\Omega} \frac{\partial u}{\partial n} \, ds.$$

This is called Green's first identity.

In this paper, we consider Sobolev-type fractional partial integrodifferential equations for  $t \in J$  of the form

$$(3) \quad \begin{aligned} & \frac{C \partial^\alpha}{\partial t^\alpha} [u(x, t) - \Delta u(x, t)] \\ & = \Delta u(x, t) + f \left( t, x, u(x, t), \int_0^t k(t, s, x, u(x, s)) ds \right), \end{aligned}$$

where  $0 < \alpha < 1$ ,  $\Omega$  is a bounded subset of  $\mathbb{R}$  with smooth boundary  $\partial\Omega$ ,  $J = [0, T]$  and  $f : J \times \Omega \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  and  $k : J \times J \times \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  are nonlinear continuous functions. The initial and Neumann boundary conditions are given by

$$(4) \quad u(x, 0) = u_0(x), \quad x \in \Omega,$$

$$(5) \quad \frac{\partial u(x, t)}{\partial n} = 0, \quad (x, t) \in \partial\Omega \times J,$$

where  $u_0(x) \in C^2(\Omega)$ . In order to establish the main result, we assume the following conditions:

(H<sub>1</sub>)  $f(t, x, u_1, u_2)$  is continuous with respect to  $u_1, u_2$ , Lebesgue measurable with respect to  $t$  and satisfies

$$\frac{1}{\text{vol}(\Omega)} \int_{\Omega} f(t, x, u_1, u_2) \, dx \leq f \left( t, \frac{\int_{\Omega} u_1(x, t) \, dx}{\text{vol}(\Omega)}, \frac{\int_{\Omega} u_2(x, t) \, dx}{\text{vol}(\Omega)} \right).$$

(H<sub>2</sub>) There exists an integrable function  $m_1 : J \rightarrow [0, \infty)$  and a constant  $L_1 > 0$  such that

$$\begin{aligned} & \|f(t, x, u_1, u_2)\| \leq m_1(t)(\|u_1\| + \|u_2\|), \\ & \int_0^T (T-s)^{\alpha-1} m_1(s) ds \leq L_1 \text{ for some } \alpha > 0. \end{aligned}$$

(H<sub>3</sub>)  $k(t, s, x, u)$  is continuous with respect to  $u$ , Lebesgue measurable with respect to  $t$  and satisfies

$$\frac{1}{\text{vol}(\Omega)} \int_{\Omega} k(t, s, x, u) \, dx \leq k \left( t, s, \frac{\int_{\Omega} u(x, t) \, dx}{\text{vol}(\Omega)} \right).$$

(H<sub>4</sub>) There exists an integrable function  $m_2(t, s) : J \times J \rightarrow [0, \infty)$  and a constant  $L_2 > 0$  such that

$$\|k(t, s, x, u)\| \leq m_2(t, s)\|u\|,$$

$$\int_0^T (T-s)^{\alpha-1} m_1(s) \int_0^s m_2(s, \tau) d\tau ds \leq L_2 \text{ for some } \alpha > 0,$$

$$L_1 + L_2 \neq \Gamma(\alpha).$$

It is easy to show that by using (1), the fractional partial integrodifferential equation (3) with (4) is equivalent to the equation

$$\begin{aligned} (6) \quad u(x, t) &= u_0(x) - u_0''(x) + \Delta u(x, t) \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \Delta u(x, s) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, x, u(x, s), v(x, s)) \, ds, \end{aligned}$$

where

$$v(x, s) = \int_0^s k(s, \tau, x, u(x, \tau)) d\tau.$$

### 3. EXISTENCE RESULT

In this section, we establish the main result of this paper. For this, we define the functions  $U(t)$  and  $V(t)$  as

$$U(t) = \frac{1}{\text{vol}(\Omega)} \int_{\Omega} u(x, t) \, dx$$

$$V(t) = \frac{1}{\text{vol}(\Omega)} \int_{\Omega} v(x, t) \, dx.$$

**Theorem 3.** *Suppose that (H<sub>1</sub>)–(H<sub>4</sub>) hold. Then there exists at least one solution for the initial value problem (3) on  $J$ .*

*Proof.* We have to prove that the initial value problem (3) has a solution if and only if the equation

$$(7) \quad U(t) = U(0) - U''(0) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, U(s), V(s)) \, ds,$$

where

$$U''(0) = \frac{1}{\text{vol}(\Omega)} \int_{\Omega} \Delta u(x, 0) \, dx = \frac{1}{\text{vol}(\Omega)} \int_{\Omega} u_0''(x) \, dx$$

has a solution.

Assume  $u(x, t)$  to be a solution of (3). Then it follows that  $u(x, t)$  is a solution of (6). Now integrating both sides of equation (6) with respect to  $x \in \Omega$ , we get

$$\begin{aligned} \int_{\Omega} u(x, t) \, dx &= \int_{\Omega} u_0(x) \, dx - \int_{\Omega} u_0''(x) \, dx + \int_{\Omega} \Delta u(x, t) \, dx \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{\Omega} \int_0^t (t-s)^{\alpha-1} \Delta u(x, s) \, ds \, dx \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{\Omega} \int_0^t (t-s)^{\alpha-1} f(s, u(x, s), v(x, s)) \, ds \, dx \end{aligned}$$

Using the Green identity (2) and the Neumann boundary condition (5) we get

$$\int_{\Omega} \Delta u(x, t) \, dx = 0.$$

By the assumptions (H<sub>1</sub>) and (H<sub>3</sub>), we have

$$(8) \quad U(t) \leq U(0) - U''(0) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, U(s), V(s)) \, ds.$$

Choose  $b \geq \frac{\|U(0)\|(L_1 + L_2)}{\Gamma(\alpha) - (L_1 + L_2)}$  and let

$$K = \{U : U \in C(J, \mathbb{R}), \|U(t) - U(0)\| \leq b\}.$$

Define the nonlinear operator  $F : C(J, \mathbb{R}) \rightarrow C(J, \mathbb{R})$  as

$$(9) \quad FU(t) = U(0) - U''(0) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, U(s), V(s)) \, ds.$$

Clearly  $U(0) \in K$  and so  $K$  is nonempty. Further  $K$  is a closed bounded convex set. Next we have to prove that the operator  $F$  maps  $K$  into itself.

Note that

$$\|FU(t) - FU(0)\| \leq \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \|f(s, U(s), V(s))\| ds.$$

Then by using  $(H_2)$  and  $(H_4)$ , we get

$$\begin{aligned} & \|FU(t) - FU(0)\| \\ & \leq \frac{1}{\Gamma(\alpha)} \int_0^t m_1(s)(t-s)^{\alpha-1} [\|U(s)\| + \|V(s)\|] ds \\ & \leq \frac{1}{\Gamma(\alpha)} (\|U(0)\| + b) \int_0^T m_1(s)(T-s)^{\alpha-1} [1 + \int_0^s m_2(s, \tau) d\tau] ds \\ & \leq \frac{L_1 + L_2}{\Gamma(\alpha)} (\|U(0)\| + b) \\ & \leq b. \end{aligned}$$

Therefore  $F$  maps  $K$  into itself. Using similar argument as in [5, 27], it is easy to show that  $F$  is completely continuous. Then by the Leray-Schauder fixed point theorem,  $F$  has a fixed point in  $K$ , which is a solution of the equation (3).  $\square$

**Remark.** Consider the Sobolev-type fractional partial integrodifferential equation with kernel of the form

$$(10) \quad \frac{C\partial^\alpha}{\partial t^\alpha} [u(x, t) - \Delta u(x, t)] = \Delta u(x, t) + \int_0^t h(t-s)\Delta u(x, s) ds + f(t, x, u(x, t)),$$

where  $f : J \times \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  and  $h : J \rightarrow \mathbb{R}$  are nonlinear continuous functions. Existence of solutions of equation (10) with Neumann boundary condition (5) can be established by using the same method as in Theorem 3.

#### 4. ABSTRACT FRACTIONAL SOBOLEV EQUATION

The Sobolev-type fractional partial integrodifferential equation (3) with (4) can be discussed in abstract setting as in [6, 14, 26] and it can be written for  $t \in J$  as

$$(11) \quad \begin{aligned} \frac{C\partial^\alpha}{\partial t^\alpha} [Bu(x, t)] &= Au(x, t) + f\left(t, x, u(x, t), \int_0^t k(t, s, x, u(x, t)) ds\right), \\ u(x, 0) &= u_0(x), \end{aligned}$$

where  $B = I - A$  and  $A = \Delta$ . Suppressing the phase variable and letting  $X$  a general Banach space, (11) can be written as abstract fractional differential equation of the form

$$(12) \quad \begin{aligned} {}^C D^\alpha [Bu(t)] &= Au(t) + f\left(t, u(t), \int_0^t k(t, s, u) ds\right), \quad t \in J, \\ u(0) &= u_0, \end{aligned}$$

where  ${}^C D^\alpha$  is the Caputo fractional derivative with  $0 < \alpha < 1$ . The operators  $A$  and  $B$  are linear with domains contained in  $X$  and ranges contained in a Banach Space  $Y$  and the operators  $A : D(A) \subset X \rightarrow Y$  and  $B : D(B) \subset X \rightarrow Y$  satisfy the following hypotheses:

- (C1)  $A$  and  $B$  are closed linear operators,
- (C2)  $D(B) \subset D(A)$  and  $B$  is bijective,
- (C3)  $B^{-1} : Y \rightarrow D(B)$  is compact,
- (C4)  $B^{-1}A : X \rightarrow D(B)$  is continuous.

The nonlinear operators  $f : J \times X \times X \rightarrow Y$  and  $k : J \times J \times X \rightarrow X$  are continuous. It is easy to prove that the equation (12) is equivalent to the integral equation

$$(13) \quad \begin{aligned} u(t) &= u_0 + \frac{1}{\Gamma(\alpha)} \int_0^t (t - s)^{\alpha-1} B^{-1} Au(s) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t - s)^{\alpha-1} B^{-1} f\left(s, u(s), \int_0^s k(s, \tau, u) d\tau\right) ds. \end{aligned}$$

By using the resolvent operators, the solvability problem for the equation (12) has been discussed in [6, 22].

### 5. EXAMPLE

Consider the Sobolev-type fractional partial integrodifferential problem of the form

$$(14) \quad \begin{aligned} \frac{\partial^{\frac{3}{4}}}{\partial t^{\frac{3}{4}}} \left[ u(t, x) - \frac{\partial^2}{\partial x^2} u(t, x) \right] &= \frac{\partial^2}{\partial x^2} u(t, x) + b(t) \sin u(t, x) \\ &+ \int_0^t a(t - s) e^{-u(s, x)} ds, \quad t > 0, \end{aligned}$$

$$(15) \quad u(x, 0) = u_0(x), \quad x \in \Omega,$$

$$(16) \quad \frac{\partial u(x, t)}{\partial n} = 0, \quad (x, t) \in \partial\Omega \times J,$$

where  $J = [0, T], \Omega = [0, \pi], u_0 \in C^2(\Omega)$  and  $a, b \in C(J)$ . Here the function  $f : J \times \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  and  $k : J^2 \times \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  are given as

$$f(t, x, u(t, x), Ku(t, x)) = b(t) \sin u(t, x) + Ku(t, x),$$

$$Ku(t, x) = \int_0^t k(t, s, x, u(s, x)) ds = \int_0^t a(t-s) e^{-u(s, x)} ds.$$

By appropriately choosing the functions  $a(t), b(t)$  [6], one can impose the conditions  $(H_1) - (H_4)$  of Theorem 3 on the functions  $f, k$ , and hence there is a function  $u \in C(J, \mathbb{R})$  which satisfies the equations (14)–(16) on  $J$ .

#### REFERENCES

- [1] A. Akilandeswari, K. Balachandran, N. Annapoorani, *Solvability of hyperbolic fractional partial differential equations*, Journal of Applied Analysis and Computation, 7 (2017), 1570-1585.
- [2] A. Akilandeswari, K. Balachandran, N. Annapoorani, *Existence of solutions of fractional partial integrodifferential equations with Neumann boundary condition*, Non-linear Functional Analysis and Applications, 22 (2017), 711-722.
- [3] A. Akilandeswari, K. Balachandran, J.J. Trujillo, M. Rivero, *On the solutions of partial integrodifferential equations of fractional order*, Tbilisi Mathematical Journal, 10 (2017), 19-29.
- [4] K. Balachandran, *An Introduction to Fractional Differential Equations*, Springer, Singapore, 2023.
- [5] K. Balachandran, *On the solutions of Sobolev type fractional partial differential equations*, The Aligarh Bulletin of Mathematics, 42 (2023), 81-94.
- [6] K. Balachandran, S. Kiruthika, *Existence of solutions of abstract fractional integrodifferential equations of Sobolev type*, Computers and Mathematics with Applications, 64 (2012), 3406-3413.
- [7] K. Balachandran, J.Y. Park, *Nonlocal Cauchy problem for Sobolev type functional integrodifferential equation*, Bulletin of the Korean Mathematical Society, 39 (2002), 561-569.
- [8] K. Balachandran, J.Y. Park, *Sobolev type integrodifferential equation with nonlocal condition in Banach spaces*, Taiwanese Journal of Mathematics, 7 (2003), 155-163.
- [9] K. Balachandran, R. Ravikumar, *Nonlocal Cauchy problem for time varying delay integrodifferential equations of Sobolev type in Banach spaces*, Tamkang Journal of Mathematics, 37 (2006), 193-205.
- [10] K. Balachandran, K. Uchiyama, *Existence of solutions of nonlinear integrodifferential equations of Sobolev type with nonlocal condition in Banach spaces*, Proceedings of the Indian Academy of Sciences (Mathematical Sciences), 110 (2000), 225-232.
- [11] K. Balachandran, J.Y. Park, M. Chandrasekaran, *Nonlocal Cauchy problem for delay integrodifferential equations of Sobolev type in Banach spaces*, Applied Mathematics Letters, 15 (2002), 845-857.

- 
- [12] K. Balachandran, D.G. Park, Y.C. Kwun, *Nonlinear integrodifferential equations of Sobolev type with nonlocal conditions in Banach spaces*, Communications of the Korean Mathematical Society, 14 (1999), 223-231.
- [13] G. Barenblat, J. Zheltor, I. Kochiva, *Basic concepts in the theory of seepage of homogeneous liquids in fissured rocks*, Journal of Applied Mathematics and Mechanics, 24 (1960), 1286-1303.
- [14] H. Brill, *A semilinear Sobolev equation in a Banach space*, Journal of Differential Equations, 24 (1977), 412-425.
- [15] S. Cetinkaya, A. Demir, *Effects of the ARA transform method for time fractional problems*, Mathematica Moravica, 26 (2022), 73-84.
- [16] L.C. Evans, *Partial Differential Equations*, American Mathematical Society, Providence, 1998.
- [17] A. Granas, J. Dugundji, *Fixed Point Theory*, Springer-Verlag, New York, 2003.
- [18] T.L. Guo, K. Zhang, *Impulsive fractional partial differential equations*, Applied Mathematics and Computation, 257 (2015), 581-590.
- [19] M.A.E. Herzallah, A.M.A. El-Sayed, D. Baleanu, *On the fractional order diffusion-wave process*, Romanian Journal of Physics, 55 (2010), 274-284.
- [20] M. Javidi, B. Ahmad, *Numerical solution of fractional partial differential equations by Laplace inversion technique*, Advances in Differential Equations, 375 (2013), 1-18.
- [21] R. Joice Nirmala, K. Balachandran, *Analysis of solutions of time fractional telegraph equation*, Journal of the Korean Society for Industrial and Applied Mathematics, 18 (2014), 209-224.
- [22] B. Kamalpriya, K. Balachandran, N. Annapoorani, *Existence results for fractional integrodifferential equations of Sobolev type with deviating arguments*, Journal of Applied Nonlinear Dynamics, 11 (2022), 57-67.
- [23] A.A. Kilbas, H.M. Srivasta, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006.
- [24] Y. Luchko, *Maximum principle for the generalized time-fractional diffusion equation*, Journal of Mathematical Analysis and Applications, 351 (2009), 218-223.
- [25] Y. Luchko, *Some uniqueness and existence results for the initial-boundary value problems for the generalized time-fractional diffusion equation*, Computers and Mathematics with Applications, 59 (2010), 1766-1772.
- [26] J.H. Lightbourne III, S.M. Rankin III, *A partial functional differential equation of Sobolev type*, Journal of Mathematical Analysis and Applications, 93 (1983), 328-337.
- [27] Z. Ouyang, *Existence and uniqueness of the solutions for a class of nonlinear fractional order partial differential equations with delay*, Computers and Mathematics with Applications, 61 (2011), 860-870.
- [28] T. Oussaeif, A. Bouziani, *Existence and uniqueness of solutions to parabolic fractional differential equations with integral conditions*, Electronic Journal of Differential Equations, 2014 (2014), 1-10.

- [29] V. Parthiban, K. Balachandran, *Solutions of systems of fractional partial differential equations*, Application and Applied Mathematics, 8 (2013), 289-304.
- [30] R.E. Showalter, *Existence and representation theorems for a semilinear Sobolev equation in Banach space*, SIAM Journal on Mathematical Analysis, 3 (1972), 527-543.

**K. BALACHANDRAN**

DEPARTMENT OF MATHEMATICS

BHARATHIAR UNIVERSITY

COIMBATORE 641 046

INDIA

*E-mail address:* kb.maths.bu@gmail.com